

ALGEBRAIC CYCLES AND MOTIVIC ITERATED INTEGRALS

II

AMIR JAFARI

ABSTRACT. This is a sequel to [FJ]. It will give a more natural framework for constructing elements in the Hopf algebra χ_F of framed mixed Tate motives according to Bloch and Křiž [BK]. This framework allows us to extend the results of [FJ] to interpret all multiple zeta values (including the divergent ones) and the multiple polylogarithms in one variable as elements of χ_F . It implies that the pro-unipotent completion of the torsor of paths on $\mathbb{P}^1 - \{0, 1, \infty\}$, is a mixed Tate motive in the sense of [BK]. Also It allows us to interpret the multiple logarithm $Li_{1, \dots, 1}(z_1, \dots, z_n)$ as an element of χ_F as long as the products of consecutive z_i 's are not 1.

CONTENTS

1. Introduction	1
2. A Combinatorial Differential Graded Algebra	4
3. The Cycle map	9
4. Hodge Realization of the Motivic Iterated Integrals	18
Appendix A. Bar Construction	22
Appendix B. Mixed Hodge-Tate Structures	23
References	26

1. INTRODUCTION

The category $\mathcal{MTM}(F)$ of mixed Tate motives over a field F is supposed to be a Tannakian category with a special invertible object $\mathbb{Q}(1)$. Let $\mathbb{Q}(-1) = \underline{\text{Hom}}(\mathbb{Q}, \mathbb{Q}(1))$. The simple objects of this category are the mutually distinct elements $\mathbb{Q}(n) = \otimes^{|n|} \mathbb{Q}(n/|n|)$ and $\mathbb{Q}(0) = \mathbb{Q}$. If we define the weight of $\mathbb{Q}(n)$ to be $-2n$, then any object of this category has a canonical increasing (weight) filtration such that $gr_{-2n+1} = 0$ and gr_{-2n} is a direct sum of $\mathbb{Q}(n)$'s. One should also have the following relation with K -theory of F :

$$\text{Ext}^i(\mathbb{Q}(0), \mathbb{Q}(n)) = gr_{\gamma}^n K_{2n-i}(F)_{\mathbb{Q}}.$$

Furthermore if F can be embedded inside the field of complex numbers $\mathbb{C}$, there should be a Hodge realization tensor functor from $\mathcal{MTM}(F)$ to the category of mixed Hodge-Tate structures. $\mathbb{Q}(1)$ will correspond to the Hodge structure of $H_2(\mathbb{P}^1)$ or equivalently $H_1(\mathbb{P}^1 \setminus \{0, \infty\})$. Also when F has a characteristic prime to a prime number ℓ and contains only finitely many ℓ^{th} power roots of 1, there should be an ℓ -adic realization tensor functor from $\mathcal{MTM}(F)$ to the category of

mixed Tate ℓ -adic representations of the absolute Galois group $G_F = \text{Gal}(\overline{F}/F)$. $\mathbb{Q}(1)$ will correspond to the action of G_F on

$$\mathbb{Q} \otimes \varprojlim \mu_{\ell^n}(\bar{F}) = \mathbb{Q}_{\ell}(1).$$

The existence of this category in this generality is still conjectural. However if the field F satisfies Beilinson-Soulé conjecture stating:

$$gr_{\gamma}^n K_{2n-i}(F)_{\mathbb{Q}} = 0 \quad \text{for } i \leq 0 \quad (\text{except when } i = n = 0)$$

such a category can be constructed as the heart of a canonical t -structure on the triangulated subcategory of mixed motives over F (constructed independently by Voevodsky, Levine and Hanamura) generated by $\mathbb{Q}(n)$'s. It is known for example thanks to Borel that a number field satisfies this conjecture.

Bloch and Křiz in [BK] proposed a different approach. Using algebraic cycles in $\square_F^n = (\mathbb{P}_F^1 \setminus \{1\})^n$ whose support meets all the faces (defined by setting some of the coordinates = 0 or ∞) properly, they constructed a graded Hopf algebra that they proposed to be the Hopf algebra of framed mixed Tate motives.

Let $\text{Cycle}^r(n)$ be admissible cycles of codimension r in $\square_F^n$. The permutation group Σ_n and the group $(\mathbb{Z}/2\mathbb{Z})^n$ act on $\square_F^n$. So we have an action of $\mathcal{G}_n = \Sigma_n \rtimes (\mathbb{Z}/2\mathbb{Z})^n$ on $\text{Cycle}^r(n)$. Let

$$\mathcal{N}^n(r) = \text{Alt}(\text{Cycle}^r(2r-n) \otimes \mathbb{Q})$$

where

$$\text{Alt} = \frac{1}{|\mathcal{G}_{2r-n}|} \sum_{\tau \in \mathcal{G}_{2r-n}} \text{sgn}(\tau) \tau.$$

The sign of an element $(\sigma, \epsilon_1, \dots, \epsilon_n)$ where $\sigma \in \Sigma_n$ and $\epsilon_i = \pm 1$ is defined to be $\text{sgn}(\sigma) \epsilon_1 \dots \epsilon_n$. The external product followed by Alt defines a product

$$\mathcal{N}^n(r) \otimes \mathcal{N}^m(s) \longrightarrow \mathcal{N}^{n+m}(r+s)$$

that is graded commutative with respect to the upper grading. Also we have a differential:

$$d : \mathcal{N}^n(r) \longrightarrow \mathcal{N}^{n+1}(r)$$

$$Z \mapsto \sum_{i=1}^{2r-n} (-1)^{i-1} (Z|_{x_i=0} - Z|_{x_i=\infty}).$$

These structures make $\mathcal{N} = \oplus \mathcal{N}^n(r)$ into a graded differential algebra (DGA) with Adams grading. The seminal works of Bloch and Levine on higher Chow groups imply that:

$$H^i(\mathcal{N}(n)^{\bullet}) = gr_{\gamma}^n K_{2n-i}(F)_{\mathbb{Q}}.$$

Hence the conjecture of Beilinson-Soulé briefed above is equivalent to the fact that $\mathcal{N}$ is cohomologically connected.

The bar construction, invented by Eilenberg-MacLane and Adams, to any DGA $\mathcal{A}$ associates a new DGA $B(\mathcal{A})$ with a coproduct. This construction is reminded in appendix A.

The proposed Hopf algebra of framed mixed Tate motives is then

$$\chi_F = H^0 B(\mathcal{N}_F).$$

In [BK] the realization morphisms from χ_F to the Hopf algebras of framed mixed Hoge-Tate structures and framed mixed ℓ -adic representations of G_F are constructed.

The category of mixed Tate motives over F is defined to be the graded co-modules over χ_F .

A rich source for constructing mixed motives is the unipotent completion of the fundamental group or the torsor of paths of smooth algebraic varieties. For any smooth variety X and $a, b \in X$, the truncated torsor of paths $\Pi^N(X, a, b)$ is defined by:

$$\frac{\mathbb{Q}[\pi(X, a, b)]}{I_a^{N+1}\mathbb{Q}[\pi(X, a, b)]}.$$

Here I_a is the augmentation ideal $\text{Ker}(\mathbb{Q}[\pi_1(X, a)] \rightarrow \mathbb{Q})$ (We are viewing X as a complex manifold by taking its points over $\mathbb{C}$, however there is an algebraic way to define these objects). According to Morgan and Hain this space has a natural mixed Hodge structure, in fact Hain and Zucker showed that $\Pi^N(X, a, b)$ defines a good unipotent variation of mixed Hodge structures on $X \times X$. Therefore even for a and b in a smooth compactification $\bar{X}$ of X and not in X , one can take limit mixed Hodge structure to define the truncated torsor of paths from a to b . It's easy to see that $gr_0\Pi^N = \mathbb{Q}(0)$. Assuming $W_{-4}H_2(X) = 0$, one can show

$$gr_{-2N}\Pi^N(X, a, b) = \otimes^N W_{-2}H_1(X)$$

(Here equality means a canonical isomorphism.) Therefore if one has non-vanishing holomorphic functions $f_1, \dots, f_N$ on X , we get a morphism:

$$f_{1*} \otimes \dots \otimes f_{N*} : \otimes^N W_{-2}H_1(X) \rightarrow \otimes^N W_{-2}H_1(\mathbb{P}^1 \setminus \{0, \infty\}) = \mathbb{Q}(N).$$

Hence we get a framed mixed Hodge structure with the canonical frame $gr_0\Pi^N = \mathbb{Q}(0)$ and the frame:

$$f_{1*} \otimes \dots \otimes f_{N*} : gr_{-2N}\Pi^N \rightarrow \mathbb{Q}(N).$$

This is the Hodge theoretic version of the Iterated integral:

$$\int_a^b \frac{df_1}{f_1} \circ \dots \circ \frac{df_N}{f_N}.$$

For convenience of the reader we recall here the definition of such integrals. Let M be a manifold and $\omega_1, \dots, \omega_n$ smooth 1-forms on M . For any path $\gamma : [0, 1] \rightarrow M$, let $\gamma^*\omega_i = f_i(t)dt$, then:

$$\int_\gamma \omega_1 \circ \dots \circ \omega_n = \int_{0 \leq t_1 \leq \dots \leq t_n \leq 1} f_1(t_1) \dots f_n(t_n) dt_1 \dots dt_n.$$

The framed mixed Hodge structure defined above, being of geometric origin, should have an upgrade to a framed mixed motive. If one assumes that for a smooth compactification $\bar{X}$ of X , $H^1(\bar{X}) = 0$, then it is shown that $\Pi^N(X, a, b)$ carries a mixed Hodge-Tate structure and therefore we expect to have an upgrade to a framed mixed Tate motive and therefore an element $\mathbb{I}_X(a; f_1, \dots, f_N; b) \in H^0 B(\mathcal{N})$.

Our goal in this article is to carry out this problem for the case $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$ completely and therefore giving motivic interpretation of the multiple zeta values:

$$\zeta(n_1, \dots, n_k) = \sum_{0 < m_1 < \dots < m_k} \frac{1}{m_1^{n_1} \dots m_k^{n_k}}$$

and more generally a motivic interpretation of the multiple polylogarithms in one variable:

$$Li_{n_1, \dots, n_k}(z) = \sum_{0 < m_1 < \dots < m_k} \frac{z^{m_k}}{m_1^{n_1} \dots m_k^{n_k}}.$$

As they can be written as iterated integrals on $\mathbb{P}^1 \setminus \{0, 1, \infty\}$:

$$\int_0^z \underbrace{\frac{dt}{1-t} \circ \frac{dt}{t} \circ \cdots \circ \frac{dt}{t}}_{n_1} \circ \cdots \circ \underbrace{\frac{dt}{1-t} \circ \frac{dt}{t} \circ \cdots \circ \frac{dt}{t}}_{n_k}.$$

Unfortunately since some of the cycle constructions are not admissible we are unable to treat the more general case for the multi-variable multiple polylogarithms:

$$Li_{n_1, \dots, n_k}(z_1, \dots, z_k) = \sum_{0 < m_1 < \dots < m_k} \frac{z_1^{m_1} \dots z_k^{m_k}}{m_1^{n_1} \dots m_k^{n_k}}$$

which also has an iterated integral definition (assuming $z_i \neq 0$):

$$\int_0^1 \underbrace{\frac{dt}{a_1-t} \circ \frac{dt}{t} \circ \cdots \circ \frac{dt}{t}}_{n_1} \circ \cdots \circ \underbrace{\frac{dt}{a_k-t} \circ \frac{dt}{t} \circ \cdots \circ \frac{dt}{t}}_{n_k}.$$

with $a_i = (z_i z_{i+1} \dots z_k)^{-1}$. However we can give a motivic interpretation of the iterated integral:

$$\int_{a_0}^{a_{n+1}} \frac{dt}{t-a_1} \circ \cdots \circ \frac{dt}{t-a_n}$$

in the following cases:

- (1) There is an element a in F such that $a_i = a_j$ for $i \neq j$ implies $a_i = a$. We call such a sequence a -generic.
- (2) For $i = 1, \dots, n$, we have $a_i \neq a_{i+1}$. We call such a sequence sequentially distinct.
- (3) There are two distinct elements α and β such that $a_i \in \{\alpha, \beta\}$ for $i = 1, \dots, n$. We call such a sequence binary.

For example the second condition allows us to give an element in $H^0 B(\mathcal{N})$ that corresponds to the multiple logarithm $Li_{1, \dots, 1}(z_1, \dots, z_k)$ with the assumption that for any $m \leq n$: $\prod_{i=m}^n z_i \neq 1$. With the more restrictive assumption on z_i 's which demands $\prod_{i \in I} z_i \neq 1$ for any subset I of $\{1, \dots, k\}$ the first condition can be used to construct an element in $H^0 B(\mathcal{N})$ that corresponds to the multi-variable multiple polylogarithm $Li_{n_1, \dots, n_k}(z_1, \dots, z_k)$.

Remark 1.1. In our joint work with Furusho, we treated the case of 0-generic sequences in the above terminology. This article is written so that it will redo the results of [FJ] in a more transparent framework and extend them to cover more cases. Therefore there are some major overlaps; the author has decided to keep them for the sake of convenience for the reader. The language of trees, which was also used by Gangl, Goncharov and Levin in [GGL], has been replaced by the combinatorial DGA introduced in section 2. It has to be stressed that the existence of this paper and [FJ] owe a great deal to [GGL].

2. A COMBINATORIAL DIFFERENTIAL GRADED ALGEBRA

Let S be a set and $\mathcal{C}_S$ the graded commutative algebra with unit, freely generated by symbols:

$$(a_0; a_1, a_2, \dots, a_n; a_{n+1})$$

of degree 1 and Adams grading n where $a_i \in S$ and $n > 0$.

Definition 2.1. A 2-cut of $A = (a_0; a_1, \dots, a_n; a_{n+1})$ is an ordered pair

$$(a_0; \dots, a_{i-1}, a_i, a_{j+1}, a_{j+2}, \dots; a_{n+1}), (a_i; a_{i+1}, \dots; a_j; a_{j+1})$$

for a choice $0 \leq i < j \leq n$, $(i, j) \neq (0, n)$.

Similarly a k -cut is an ordered k -tuple obtained from a $(k-1)$ -cut $A_1, \dots, A_{k-1}$ by replacing one of the A_i 's with a 2-cut A'_i and A''_i . We also define a 1-cut of A to be just A . Note that a cut should be regarded in terms of the indices and not the elements. So two 2-cuts that correspond to two different choices for $i < j$ are considered to be different even if their realized pairs are identical.

Define the differential on $\mathcal{C}_S$ for a generator $A = (a_0; a_1, \dots, a_n; a_{n+1})$ to be

$$dA = - \sum A' A''$$

where the sum is over all 2-cuts of A .

Lemma 2.2. $d^2 = 0$.

Proof. Let $A' = (a_0, \dots, a_i, a_{j+1}, \dots, a_{n+1})$, $A'' = (a_i; \dots; a_{j+1})$ be a 2-cut of A . A 2-cut A'_1, A'_2 of A' is said to be joint with A'' if A'_2, A'' is a 2-cut of a subsequence $A_3 = (a_k; a_{k+1}, \dots; a_{l+1})$ of A . otherwise we call (A'_1, A'_2) disjoint from A'' . Now

$$ddA = \sum A'_1 A'_2 A'' - \sum A' A'_1 A''_2.$$

The terms in this sum cancel each other as follows. If A'_1, A'_2 is disjoint from A'' then can consider removing A'_2 from A first and then remove A'' from the remaining of A , we will get $A'_1 A'' A'_2$ that cancels $A'_1 A'_2 A''$ due to graded commutativity of $\mathcal{C}_S$. If the cut is joint and A'_2, A'' is a 2-cut of A_3 as above, we can remove A_3 from A first and we get the term $A'_1 A_3$ and then cut A_3 into A'_2 and A'' , this cancels $A'_1 A'_2 A''$ since it appears with a negative sum in the formula above for d^2 . A similar argument for terms of the form $A' A'_1 A'_2$ finishes the proof. $\square$

Definition 2.3. Associated to a generator $A = (a_0; a_1, \dots, a_n; a_{n+1})$ define $T(A) \in B(\mathcal{C}_S)^0$ by:

$$T(A) = \sum [A_1 | \dots | A_k]$$

where the sum is over all cuts of A (including 1-cut).

Lemma 2.4. For $A = (a_0; a_1, \dots, a_n; a_{n+1})$, $d(T(A)) = 0$. Therefore $T(A) \in H^0 B(\mathcal{C}_S)(n) \subseteq B(\mathcal{C}_S)^0$.

Proof. This follows once we observe:

$$d_{ext}(\sum_{k\text{-cuts}} [A_1 | \dots | A_k]) + d_{int}(\sum_{(k+1)\text{-cuts}} [A_1 | \dots | A_{k+1}]) = 0.$$

$\square$

We will need the following lemma in section 4.

Lemma 2.5. For $A = (a_0; a_1, \dots, a_n; a_{n+1})$ let $\mathcal{C}_A$ be the sub DGA of $\mathcal{C}_S$ generated by $A_1, \dots, A_k$ for all cuts of A . Then $H^0 B(\mathcal{C}_A)$ is generated by $T(A_i)$'s as an algebra.

Proof. It can be proved via a counting argument. Details are left to the reader. $\square$

Proposition 2.6. *The coproduct of $T(A)$ is given by:*

$$1 \otimes T(A) + T(A) \otimes 1 + \sum T(A_1) \otimes T(A_2) \cdots T(A_k)$$

where the sum is over all elementary k -cuts of A for $k > 1$. An elementary k -cut $A_1, \dots, A_k$ is a cut with the property that each A_i for $i > 1$ is formed of consecutive elements, i.e. for some $l \geq 0$ and $m > 1$ it is of the form $(a_l; a_{l+1}, \dots, a_{l+m-1}; a_{l+m})$.

Remark 2.7. This formula is analogous to Goncharov's formula for the coproduct of the framed mixed Hodge-Tate structure associated to iterated integrals in [G], theorem 6.1.

Proof. By definition of coproduct

$$\Delta T(A) = 1 \otimes T(A) + T(A) \otimes 1 + \sum [B_1 | \dots | B_m] \otimes [B_{m+1} | \dots | B_l]$$

where the sum is over all l -cuts of A with $l > 1$ and $0 < m < l$. Fix an elementary k -cut $c = (A_1, \dots, A_k)$ of A . A term $[B_1 | \dots | B_m] \otimes [B_{m+1} | \dots | B_l]$ is said to be of type c if $B_1, \dots, B_m$ is an m -cut of A_1 . This relation partitions the terms of the above sum into groups indexed by elementary cuts of A . If $[B_1 | \dots | B_m] \otimes [B_{m+1} | \dots | B_l]$ is of type $c = (A_1, \dots, A_k)$ then obviously $B_{m+1}, \dots, B_l$ is a shuffle permutation of cuts of $A_2, \dots, A_k$ and therefore the sum of terms of type c gives $T(A_1) \otimes T(A_2) \cdots T(A_k)$. $\square$

Let $\tilde{\mathcal{C}}_S$ be a quotient of $\mathcal{C}_S$ by the following two types of relations:

- (1) $(a_0; a_1, \dots, a_n; a_{n+1}) + (a_{n+1}, a_1, \dots, a_n; a_{n+2}) = (a_0; a_1, \dots, a_n; a_{n+2})$.
- (2) $\sum_{\sigma \in \Sigma_{n,m}} (a_0; a_{\sigma(1)}, \dots, a_{\sigma(n+m)}; a_{n+m+1}) = 0$.
- (3) $(a_0; a_1, \dots, a_n; a_{n+1}) = (-1)^n (a_{n+1}; a_n, \dots, a_1; a_0)$.

Here $\Sigma_{n,m}$ is the subgroup of (n, m) shuffles in Σ_{n+m} consisting of those elements σ such that: $\sigma^{-1}(1) < \dots < \sigma^{-1}(n)$ and $\sigma^{-1}(n+1) < \dots < \sigma^{-1}(n+m)$. The ideal generated by these elements is both homogeneous and is stable under the differential and therefore $\tilde{\mathcal{C}}_S$ is a DGA. Let $\tilde{T}(A) \in B(\tilde{\mathcal{C}}_S)$ be the image of $T(A)$.

Proposition 2.8. *We have the following relations for $\tilde{T}(A)$:*

$$\begin{aligned} \sum_{i=0}^n \tilde{T}(a_0; a_1, \dots, a_i; a_{n+1}) \tilde{T}(a_{n+1}; a_{i+1}, \dots, a_n; a_{n+2}) &= \tilde{T}(a_0; a_1, \dots, a_n; a_{n+2}) \\ \tilde{T}(a_0; a_1, \dots, a_n; a_{n+m+1}) \tilde{T}(a_0; a_{n+1}, \dots, a_{n+m}; a_{n+m+1}) &= \\ &\sum_{\sigma \in \Sigma_{n,m}} \tilde{T}(a_0; a_{\sigma(1)}, \dots, a_{\sigma(n+m)}; a_{n+m+1}) \\ \tilde{T}(a_0; a_1, \dots, a_n; a_{n+1}) &= (-1)^n \tilde{T}(a_{n+1}; a_n, \dots, a_1; a_0) \end{aligned}$$

For the first relation, we let $T(a_0; a_{n+1}) = T(a_{n+1}; a_{n+2}) = 1$.

Remark 2.9. These formulae are analogous to the path composition and shuffle product relation for iterated integrals.

Proof. To prove the first relation, let $A = (a_0; \dots; a_{n+2})$, $B^i = (a_0; a_1, \dots, a_i; a_{n+1})$ and $C^i = (a_{n+1}; a_{i+1}, \dots, a_n; a_{n+2})$. Let $A_1, \dots, A_m$ be a cut of A such that $A_1 = A_{i_1}, \dots, A_{i_k}$ end with a_{n+2} . One can replace $A_{i_j} = (a_l; \dots; a_{n+2})$ by the sum $A'_{i_j} + A''_{i_j}$ where $A'_{i_j} = (a_l; \dots; a_{n+1})$ and $A''_{i_j} = (a_{n+1}; \dots; a_{n+2})$. Therefore $[A_1 | \dots | A_k]$ can be written as:

$$[A'_{i_1} | \dots | A'_{i_{k-1}} | \dots | A'_{i_k} | A_{i_k+1} | \dots | A_m] +$$

$$\sum_{l=1}^k [A'_{i_1} | \dots | \dots | A'_{i_{l-1}} | \dots | A''_{i_l} | A_{i_{l+1}} | \dots | A_m]$$

If $A''_{i_l} = (a_{n+1}; a_{i+1}, \dots; a_{n+2})$, this element belongs to a term in $T(B^i)T(C^i)$. This way we get a 1-1 correspondence between the terms in $T(A)$ and the terms in $\sum_{i=0}^n T(B^i)T(C^i)$.

To prove the second relation, observe that the right hand side can be written as:

$$\sum_k [A_1 | A_2 | \dots | A_k]$$

where $(A_1, \dots, A_k)$ runs through all k -cuts of $A = (a_0; a_{\sigma(1)}, \dots, a_{\sigma(n+m)}, a_{n+m+1})$ for some $\sigma \in \Sigma_{n,m}$. If $A_i = (a_{i_0}; a_{i_1}, \dots, a_{i_l}; a_{i_{l+1}})$ we let the support of A_i be the set $\{i_1, \dots, i_l\}$. We say A_i is mixed if its support has both elements $\leq n$ and $> n$. We say A_i is sequential if there is a $\sigma \in \Sigma_{n,m}$ and a subsequence A'_i , such that (A'_i, A_i) (in this order) is a 2-cut of $(a_0; a_{\sigma(1)}, \dots, a_{\sigma(n+m)}, a_{n+m+1})$. Note that the last piece A_k of a k -cut of $(a_0; a_{\sigma(1)}, \dots, a_{\sigma(n+m)}, a_{n+m+1})$ is always sequential. If A_k is mixed and we sum over all shuffles of its support $\{i_1, \dots, i_l\}$ of elements $\leq n$ and $> n$, leaving all other indices fixed, we find out that the terms $[A_1 | \dots | A_k]$ corresponding to this cut, will cancel. So A_k is pure, i.e.

$$A_k = (a_p; a_i, a_{i+1}, \dots, a_{i+j-1}; a_q)$$

where $(i, i+1, \dots, i+j-1)$ are either all $\leq n$ or all $> n$. Assume they are all $\leq n$. I claim that we can also assume without loss of generality that $p = i-1$ and $q = i+j$ if $i+j \leq n$ or $q = m+n+1$ if $i+j = n+1$, without changing the other pieces $A_1, \dots, A_{k-1}$. We give the arguments for p , the other index has a similar reasoning. If $p \neq i-1$ then it has to be bigger than n so we can swap a_p with $a_i, \dots, a_{i+j-1}$. Now the index before p is either $i-1$ or $p-1$. In the first case by removing $a_i, \dots, a_{i+j-1}$ and doing the same cuts as before from the new sequence we get:

$$A_1, \dots, A_{k-1}, (a_{i-1}; a_i, \dots, a_{i+j-1}; a_p)$$

But by the first type relations we have:

$$(a_{i-1}; a_i, \dots, a_{i+j-1}; a_p) + (a_p, a_i, \dots, a_{i+j-1}; a_q) = (a_{i-1}; a_i, \dots, a_{i+j-1}; a_q)$$

and so we have changed p to $i-1$ without changing the rest of the cut. In the second case we can replace A_k with $(a_{p-1}; a_i, \dots, a_{i+j-1}; a_q)$ and we can repeat the argument by decreasing p until we get to $i-1$. Now we pick A_{k-1} ; a similar argument shows that if its support is mixed then after summing over all the shuffles fixing the rest of the indices it will be canceled. And if it is pure, we can assume that its two ends are also in the same range. Continuing this way we can replace the terms in the right hand side with the terms in the left hand side. A neater argument can be given by an induction on $n+m$.

The last relation is easy, and its proof is omitted. $\square$

Remark 2.10. It is interesting to observe that for proving the path composition we only used the first type relations and naively one expects that for the shuffle relation we only need the second type relations. However our proof uses both relations in order to prove the shuffle relation.

We need the following lemma to deal with divergent iterated integrals

$$\int_{a_0}^{a_{n+1}} \frac{dt}{t-a_1} \circ \cdots \circ \frac{dt}{t-a_n}$$

where $a_0 = a_1$ or $a_n = a_{n+1}$.

Lemma 2.11. *The image of $\tilde{T} : \tilde{\mathcal{C}}_S \rightarrow B(\mathcal{C}_S)^0$ is generated by $\tilde{T}(a_0; a_1, \dots, a_n; a_{n+1})$ with $a_0 \neq a_1$ and $a_n \neq a_{n+1}$ (by analogy we call these elements convergent) and $\tilde{T}(b_0; b_0; b_1)$.*

Proof. Note that the shuffle relation implies:

$$\tilde{T}(a_0; \underbrace{a_1, a_1, \dots, a_1}_n; a_{n+1}) = \frac{1}{n!} (\tilde{T}(a_0; a_1; a_{n+1}))^n$$

We prove the lemma for the following type elements by an induction on $i + j$:

$$\tilde{T}(a_0; \underbrace{a_0, \dots, a_0}_i, a_{i+1}, \dots, a_{n-j}, \underbrace{a_{n+1}, \dots, a_{n+1}}_j; a_{n+1})$$

where $a_{i+1} \neq a_0$ and $a_{n-j} \neq a_{n+1}$, we may also assume $a_0 \neq a_{n+1}$. Consider the product:

$$\tilde{T}(a_0; \underbrace{a_0, \dots, a_0}_i; a_{n+1}) \tilde{T}(a_0; a_{i+1}, \dots, a_{n-j}; a_{n+1}) \tilde{T}(a_0; \underbrace{a_{n+1}, \dots, a_{n+1}}_j; a_{n+1})$$

By shuffle relation it is a linear sum of elements of the form $\tilde{T}(A)$, including the above term. Now notice that other than that term all the other terms by induction hypothesis can be written in terms of the convergent elements. $\square$

Let us define an ind-pro-object $\Pi^C(S, a, b)$ for $a, b \in S'$ with $S \subseteq S'$, in the category of graded $H^0 B(\mathcal{C}_{S, S'})$ comodule. Here $\mathcal{C}_{S, S'}$ is the sub algebra of $\mathcal{C}_{S'}$ generated by

$$(a_0; a_1, \dots, a_n; a_{n+1}) \quad \text{such that} \quad a_i \in S \quad i = 1, \dots, n.$$

When S is a finite subset of a field F , this will be a combinatorial analogue of the motivic torsor of paths $\Pi^{uni}(\mathbb{A}^1 \setminus S, a, b)$.

A graded $H^0 B(\mathcal{C}_{S, S'})$ comodule is a graded $\mathbb{Q}$ -vector space V and a graded linear map $\nu : V \rightarrow V \otimes H^0 B(\mathcal{C}_{S, S'})$ such that:

$$(id \otimes \Delta) \circ \nu = (\nu \otimes id) \circ \nu$$

$$(id \otimes \epsilon) \circ \nu = id$$

Definition 2.12. The ind-pro-object $\Pi^C(S, a, b)$ is the graded $\mathbb{Q}$ -vector spaces of non-commuting power series with variables X_s , one for each element of S , $\mathbb{Q}\langle\langle X_s \rangle\rangle_{s \in S}$. This is an ind-object with respect to S and a pro-object with respect to the polynomial degree or Adams grading. The grading is obtained by giving each X_s weight 2. For a sequence $A = (a; a_1, \dots, a_n; b) \in \mathcal{C}_{S, S'}$ let $X_A = X_{a_1} \cdots X_{a_n}$. Define:

$$\begin{aligned} \nu : \Pi^C(S, a, b) &\longrightarrow \Pi^C(S, a, b) \otimes B(\mathcal{C}_{S, S'})^0 \\ X_A &\mapsto \sum X_{A_1} \otimes [A_2] \cdots [A_k] \end{aligned}$$

where we sum over all cuts of A

If we let the graded Adams piece $H^0B(\mathcal{C}_{S,S'})(r)$ to have weight $2r$, it is readily checked that above map is a graded linear map.

Proposition 2.13. $\Pi^C(S, a, b)$ is an ind-pro-object in the category of graded $H^0B(\mathcal{C}_{S,S'})$ comodules. In particular ν factors through $\Pi^C(S, a, b) \otimes H^0B(\mathcal{C}_{S,S'})$. If we make the construction with $\tilde{\mathcal{C}}_S$, instead of $\mathcal{C}_S$, there is a natural (path composition) morphism:

$$\Pi^C(S, a, b) \otimes \Pi^C(S, b, c) \longrightarrow \Pi^C(S, a, c)$$

in the category of ind-pro graded $H^0B(\tilde{\mathcal{C}}_{S,S'})$ comodules.

Proof. The coaction can be written as:

$$X_A \mapsto \sum X_{A_1} \otimes T(A_2) \cdots T(A_k)$$

where we sum over all elementary cuts of A . The rest follows directly from coproduct formula and path composition. $\square$

3. THE CYCLE MAP

Let F be a field. Our goal is to construct a natural map of DGA's:

$$\rho : \mathcal{C}_F \longrightarrow \mathcal{N}_F.$$

This induces a map

$$\rho : H^0B(\mathcal{C}_F) \longrightarrow H^0B(\mathcal{N}_F)$$

Definition 3.1. For $A = (a_0; a_1, \dots, a_n; a_{n+1}) \in \mathcal{C}_F$, relative to ρ we define the motivic integral $\mathbb{I}_\rho(A) \in H^0B(\mathcal{N}_F)$ by $\rho(\tilde{T}(A))$.

Remark 3.2. Since $\int_a^a \omega = 0$, we expect to have $\rho(a_0; a_1, \dots, a_n; a_0) = 0$. Therefore we will replace $\mathcal{C}_F$ with its quotient by the ideal generated by $(a_0; a_1, \dots, a_n; a_0)$.

We confine ourself to construct ρ on one of the following three sub-DGA's of $\mathcal{C}_F$:

- (1) The sub-DGA $\mathcal{C}_F^a$ of a -generic elements is generated by:

$$(a_0; a_1, \dots, a_n; a_{n+1}) \text{ such that } a_i = a_j \text{ and } i \neq j \Rightarrow a_i = a.$$

- (2) The sub-DGA $\mathcal{C}_F^1$ of sequentially distinct elements¹ is generated by:

$$(a_0; a_1, \dots, a_n; a_{n+1}) \text{ such that } a_i \neq a_{i+1} \text{ for } i = 1, \dots, n.$$

- (3) The sub-DGA $\mathcal{C}_F^2$ of binary elements is generated by:

$$(a_0; a_1, \dots, a_n; a_{n+1}) \text{ such that } a_i \in \{0, 1\} \text{ for } i = 1, \dots, n.$$

The technical definition that follows, is needed for the Hodge realization. We want to construct cycles over the field $F(s_1, \dots, s_k)$. For fixed base points a_0 and a_{n+1} , there is a (partially defined) map:

$$\delta : \mathcal{N}_{F(s_1, \dots, s_k)} \longrightarrow \mathcal{N}_{F(s_1, \dots, s_{k-1})}$$

$$\begin{aligned} (\delta Z)(s_1, \dots, s_{k-1}) &= Z(a_0, s_1, \dots, s_{k-1}) - Z(s_1, s_1, \dots, s_{k-1}) + \dots \\ &\quad + (-1)^{k-1} Z(s_1, \dots, s_{k-2}, s_{k-1}, s_{k-1}) + (-1)^k Z(s_1, \dots, s_{k-1}, a_{n+1}) \end{aligned}$$

We say partially defined because the terms of this sum should be admissible. The cycles that we construct have very simple (in fact linear) coefficients in terms of s_i 's and so these specializations are sensible.

¹Note that if A is sequentially distinct and A', A'' is a 2-cut, then A' might not be sequentially distinct, however in this case $A'' = (a; \dots; a)$, an element that is set to be zero, so $\mathcal{C}_F^1$ is really a sub-DGA.

Definition 3.3. An integration theory on a sub-DGA $\mathcal{C}'_F$ of $\mathcal{C}_F$, is a sequence of algebra morphism (not DGA) for $k \geq 1$:

$$\rho_k : \mathcal{C}'_F \longrightarrow \mathcal{N}_{F(s_1, \dots, s_k)}[k-1]$$

where the bracket denotes a shift in the cohomological degree. i.e. if $A \in \mathcal{C}'_F(n)$ then $\rho_k(A) \in \mathcal{N}_{F(s_1, \dots, s_k)}^k(n)$, is a codimension n cycle inside $\square^{2n-k}$. Let $A = (a_0; a_1, \dots, a_n; a_{n+1}) \in \mathcal{C}'_F$. The following properties should hold:

- (1) $\rho_k(A) = 0$ for $k > n$.
- (2) $\rho_n(A) = \text{Alt}(c_1(s_1 - a_1), \dots, c_n(s_n - a_n))^2$ for some non-zero constants c_i 's that might depend on A .
- (3) The differential of ρ_k is:

$$d\rho_k(A) = -\delta\rho_{k+1}(A) + (-1)^k \sum_{\text{2-cuts of } A} \rho_k(A')\delta\rho_1(A'')$$

Implicit in this definition is admissibility of the cycles $\rho_k(A)$ and $\delta\rho_k(A)$ for $(k, n) \neq (1, 1)$. The case $\delta\rho_1(a_0; a_1; a_2)$ needs special definition when $a_0 = a_1$ or $a_1 = a_2$.

Lemma 3.4. *The differential of $\delta\rho_k(A)$ is given by:*

$$d(\delta\rho_k(A)) = (-1)^k \sum_{\text{2-cut of } A} \delta\rho_k(A')\delta\rho_1(A'').$$

Proof. Take boundary of both sides of equation in condition (3). Use the evident facts that $\delta d = d\delta$ and $\delta^2 = 0$ and $\delta(ab) = (\delta a)(\delta b)$. $\square$

Remark 3.5. According to lemma 3.4 the map $\rho := \delta\rho_1 : \mathcal{C}'_F \longrightarrow \mathcal{N}$ is a morphism of DGA's. So if we have an integration theory we have an associated motivic integral in $H^0B(\mathcal{N})$. In the next section we show that it has the right Hodge realization, regardless of the integration theory used.

Definition 3.6. An integration theory is called base point free, if $\rho_k(a_0; \dots; a_{n+1})$ is independent of a_0 and a_{n+1} . So for such theories we sometimes use the notation $\rho_k(a_1, \dots, a_n)$. An integration theory is called permuting if it is base point free and for all $n, m > 0$:

$$\sum_{\sigma \in \Sigma_{n,m}} \rho_1(a_{\sigma(1)}, \dots, a_{\sigma(n+m)}) = 0$$

$$\rho_1(a_1, a_2, \dots, a_n) = (-1)^{n-1} \rho_1(a_n, a_{n-1}, \dots, a_1)$$

If we have a permuting theory the corresponding $\rho : \mathcal{C}_F \longrightarrow \mathcal{N}$ factors through $\tilde{\mathcal{C}}_F$, and hence its motivic integrals $\mathbb{I}(A)$ satisfy the shuffle, path composition and inversion relations similar to those in proposition 2.8.

The construction of an integration theory is achieved inductively.

$$\hat{\rho}_1(a_1)(s_1) = \begin{cases} \text{Alt}(1 - \frac{s_1}{a_1}) & \text{if } a_1 \neq 0 \\ \text{Alt}(s_1) & \text{if } a_1 = 0 \end{cases}$$

²All is needed for the next section is: $\int_{\eta_1} \frac{dz_1}{z_1} \wedge \dots \wedge \frac{dz_n}{z_n} = \int_{\gamma} \frac{dt}{t-a_1} \circ \dots \circ \frac{dt}{t-a_n}$ where η_1 is the topological cycle obtained from $\rho_1(A)$ by replacing s_i 's with $\gamma(s_i)$'s with $0 \leq s_1 \dots \leq s_n \leq 1$ for a path γ from a_0 to a_{n+1} in $\mathbb{C} - \{a_1, \dots, a_n\}$.

We also let :

$$Sp|_{s_1=a_0}\hat{\rho}_1(a_1)(s_1) = \begin{cases} \hat{\rho}_1(a_1)(a_0) & \text{if } a_0 \neq a_1 \\ \hat{\rho}_1(a_1)(1+a_0) & \text{if } a_0 = a_1 \end{cases}$$

This specialization is used to regularize divergent integrals. They reflect the fact that with the standard tangential base point at 0 one should define $\int_0^1 \frac{dz}{z}$ to be zero. For any integration theory the most important part is a definition for $\rho_1(A)$, for $A = (a_0; a_1, \dots, a_n; a_{n+1})$. In fact if we let $A_i = (a_0; a_1, \dots, a_i; a_{i+1})$ and $A'_i = (a_i; a_{i+1}, \dots, a_n; a_{n+1})$, we can define ρ_k inductively:

$$\rho_k(A)(s_1, \dots, s_k) = \sum_{i=1}^{n-k+1} \rho_1(A_i)(s_1) \cdot \rho_{k-1}(A'_i)(s_2, \dots, s_k)$$

Lemma 3.7. *If $\rho_1(A)$ satisfies $d\rho_1(A) = -\delta\rho_2(A) - \sum \rho_1(A')\delta\rho_1(A'')$, with ρ_k 's defined as above in terms of ρ_1 for all A of depth $< n$. Then we have*

$$d\rho_k(A) = -\delta\rho_{k+1}(A) + (-1)^k \sum \rho_k(A')\delta\rho_1(A'')$$

for all A of depth $< n+k-1$.

Proof. By definition:

$$d\rho_k(A) = \sum_{i=1}^{m-k+1} d\rho_1(A_i) \cdot \rho_{k-1}(A'_i) - \rho_1(A_i) \cdot d\rho_{k-1}(A'_i)$$

where m is the depth of A , if $m < n+k-1$ then $i < n$ so we can use:

$$d\rho_1(A_i) = -\delta\rho_2(A_i) - \sum_{\text{2-cuts of } A_i} \rho_1(A_{i1})\delta\rho_1(A_{i2})$$

and inductively:

$$d\rho_{k-1}(A'_i) = -\delta\rho_k(A'_i) - \sum_{\text{2-cuts of } A'_i} \rho_{k-1}(A'_{i1})\delta\rho_1(A'_{i2})$$

so $d\rho_k(A)$ is the sum of the following two terms:

$$\begin{aligned} & \sum -\delta\rho_2(A_i) \cdot \rho_{k-1}(A'_i) + \rho_1(A_i) \cdot \delta\rho_k(A'_i) \\ & (-1)^k \sum_i \left(\sum \rho_1(A_{i1})\rho_{k-1}(A'_i)\delta\rho_1(A_{i2}) + \sum \rho_1(A_i)\rho_{k-1}(A'_{i1})\delta\rho_1(A'_{i2}) \right) \end{aligned}$$

One can check that by definition of ρ_k and fixing A_{i2} or A'_{i2} and letting i vary that the second sum is just

$$(-1)^k \sum_{\text{2-cuts of } A} \rho_k(A')\delta\rho_1(A'')$$

For the first sum if we neglect the two terms $\rho_2(A_i)(s, a_{i+1})$ and $\rho_k(A'_i)(a_i, s_1, \dots, s_{k-1})$ we get $-\delta\rho_{k+1}(A)$, now one sees immediately that these two terms when summed over i cancel each other. $\square$

Let

$$\rho_1(a_0; a_1; a_2)(s) = \begin{cases} \hat{\rho}_1(a_1 - a)(s - a) & \text{for } a\text{-generic case} \\ \hat{\rho}_1(a_1 - a_0)(s - a_0) & \text{for sequentially distinct case} \\ \hat{\rho}_1(a_1)(s) & \text{for binary case} \end{cases}$$

Note that the first and last case are base point independent. These come with their specialization coming from $\hat{\rho}_1$. We set:

$$\rho(a_0; a_1; a_2) := \delta\rho_1 := Sp|_{s=a_0}\rho_1(a_0; a_1; a_2)(s) - Sp|_{s=a_2}\rho_1(a_0; a_1; a_2)(s).$$

To define $\rho_1(A)$ for $n > 1$, we assume that we have defined ρ_1 for all elements of depth $< n$ so we have a definition of ρ_k for all elements of depth $< n + k - 1$. For the a -generic and the sequentially distinct case, let:

$$\rho_1(A)(s) = \text{Alt}\left(\frac{s-t}{c-t}, \rho_2(t, t)\right)$$

where $c = a$ in the a -generic case and $c = a_0$ for the sequentially distinct case. Here we let t vary. In fact inductively one sees that $\rho_1(A)$ has $n - 1$ of these internal variables, i.e., the way ρ_1 is defined gives us a (formal) linear combination of rational functions over $F(s)$:

$$\phi_i : (\mathbb{P}^1)^{n-1} \longrightarrow (\mathbb{P}^1)^{2n-1}.$$

The cycle is the locus of these functions, intersected with $\square^{2n-1}$. More precisely we take the pull-back of these function under the inclusion $\square^{2n-1} \hookrightarrow (\mathbb{P}^1)^{2n-1}$ to get

$$\phi'_i : X_i \longrightarrow \square^{2n-1} \quad X_i = (\mathbb{P}^1)^{n-1} \times_{(\mathbb{P}^1)^{2n-1}} \square^{2n-1}$$

These maps are proper so we have push forward ϕ'_{i*} for cycles and the cycle is:

$$\rho_1(A) = \sum_i \phi'_{i*}(X_i).$$

The definition for the binary case is more involved. One sees immediately that ρ_1 is base point dependent (in fact just the initial base point) for the sequentially distinct case and it is base point independent for the a -generic case.

Proposition 3.8. *The morphisms ρ_k define an integration theory for a -generic and sequentially distinct sub algebras. For a -generic the theory is permuting, hence we have shuffle, path composition and inversion relations for $\mathbb{L}_\rho$.*

Proof. Assume $\rho_1(A)$ is admissible and satisfies

$$d\rho_1(A) = -\delta\rho_2(A) - \sum \rho_1(A')\delta\rho_1(A'').$$

for A of depth $< n$. By lemma 3.7 for A of depth n we have:

$$d\rho_2(A) = -\delta\rho_3(A) + \sum \rho_2(A')\delta\rho_1(A'')$$

but if we specialize at (t, t) we can replace $-\delta\rho_3(t, t)$ with $-\rho_3(a_0, t, t) + \rho_3(t, t, a_{n+1})$. Assuming the admissibility, differential of $\rho_1(A) = (\frac{s-t}{c-t}, \rho_2(A)(t, t))$ is:

$$\rho_2(s, s) + (\frac{s-t}{c-t}, \rho_3(a_0, t, t)) - (\frac{s-t}{c-t}, \rho_3(t, t, a_{n+1})) - (\frac{s-t}{c-t}, \sum \rho_2(A')(t, t)\delta\rho_1(A''))$$

The first term is by $t = s$ for the first coordinate, also when $t = c$ the second factor becomes empty, This is because $\rho_2(A)(c, c)$ has at least one coordinate which is 1. The change in sign happens since the coordinates of ρ_2 are shifted by 1 in the definition of ρ_1 . Now if we use the definitions we see that this is exactly the formula we wanted to prove for $d\rho_1$. It is left to check that $\rho_1(A)$ is admissible, i.e. its intersection with any face $\mathcal{F}$ of $\square^{2n-1}$ is either empty or is of codimension n . First let's prove inductively that the intersection of $\rho_1 \cap \mathcal{F}$ is empty if we set $s = c$ (c is a_0 for sequentially distinct and is a for the generic). Recall that

$\rho_1(A)(s) = \text{Alt}(\frac{s-t}{c-t}, \rho_2(A)(t, t))$. It is enough to prove this claim before alteration. If $\mathcal{F}$ contains $z_1 = \infty$ then we have to set $t = c$ in $\rho_2(A)(t, t)$ which by induction will give empty. If the face doesn't contain $z_1 = 0$ then $\frac{s-t}{c-t}$ survives after intersection and if we let $s = c$ it is 1 so the intersection with $\square^m$ is empty. Finally if it does have $z_1 = 0$ we have to take the intersection $\rho_2(s, s)$ which is $\sum \rho_1(A_i)(s) \rho_1(A'_i)(s)$ and by induction this becomes empty if we set $s = c$. Next we show that the intersection of ρ_1 with any face $z_i = \infty$ is empty. We did the case $z_1 = \infty$ before. For other cases we have to take the intersection of $\rho_2(t, t)$ with an infinite face, if this is non-empty by induction we need to have $t = \infty$ which makes $\frac{s-t}{c-t}$ to be 1. The best way to prove the admissibility in general is to use the language of trees. We refer the reader to [FJ] or [GGL] for this. For a rooted tree T (not necessarily binary) with leaves labeled with $a_1, \dots, a_n$ and E edges, define a subvariety of $\square^E$ by:

$$\rho_1^T(A)(s) = (\frac{s-t}{c-t}, \rho_1^{T_1}(\text{supp}(T_1))(t), \dots, \rho_1^{T_k}(\text{supp}(T_k))(t))$$

where $T_1, T_2, \dots, T_k$ are the children sub-trees of the root from left to right. And if T_i has leaves labeled with $a_j, a_{j+1}, \dots, a_m$ then $\text{supp}(T_i) = (a_{j-1}; a_j, \dots, a_m; a_{m+1})$. It is observed readily that:

$$\rho_1(A)(s) = \sum \rho_1^T(A)(s)$$

where we sum over the binary trees with leaves labeled at $a_1, \dots, a_n$. Now note that the intersection of $\rho^T(A)$ with $z_i = 0$ where the coordinate corresponds to an internal edge e of T gives $\rho_1^{T'}(s)$ where T' is obtained from T by contracting e . And its intersection with $z_i = 0$ corresponding to a leaf labeled at a_i and with its sibling sub-trees $T_1, \dots, T_k$ corresponds to

$$\pm \rho_1^{T'}(\text{supp}(T'))(s) \rho^{T_1}(\text{supp}(T_1))(a_i) \dots \rho_1^{T_k}(\text{supp}(T_k))(a_i).$$

where T' is obtained from T by contracting the leaf with label a_i and cutting off all its siblings and labeling the new made leaf by a_i . So if we want to use induction on the number of edges, we only need to show that the case where one of the T_i 's above is a leaf with label at a_i will give an empty cycle. In the case of a -generic we need to have $a_i = a$ and in this case at least one other T_i gives 1 if we replace its s variable with a (note that if all T_i 's are leaves with labels at a the alteration will kill this term.) for the sequentially distinct case, the sibling T_1 to the right of the contracted leaf at a_i has its support of the form $(a_i; a_{i+1}, \dots, a_{m+1})$ and hence $\rho_1^{T_1}(\text{supp}(T_1))(a_i)$ is empty inside $\square^l$. This finishes the admissibility condition. The fact that the integration theory on a -generic elements is permuting is left to the reader. $\square$

Now we deal with the binary case. We have to define $\rho_1(a_0; a_1, \dots, a_n; a_{n+1})$, but since the theory that we will construct is base point independent, we write $\rho_1(A)$ for $A = (a_1, \dots, a_n)$. We need axillary cycles $\rho_1^\epsilon(A)$ and $\rho_1^\epsilon(A_i|A'_i)$ for $\epsilon = 0$ or 1 and $A_i = (a_1, \dots, a_i)$, $A'_i = (a_{i+1}, \dots, a_n)$. The definitions are inductive. First we give the special and simple case of $n = 2$:

$$\rho_1^\epsilon(A) = \rho_1^\epsilon(A_1|A'_1) = \frac{1}{2} \left(\frac{s-t}{\epsilon-t}, \rho_1(a_1)(t), \rho_1(a_2)(t) \right)$$

Note that by alteration this term is zero when $a_1 = a_2$. To make our formulation more compact, we also set:

$$\rho_1^\epsilon(a)(s) = |\epsilon - a| \rho_1(a)(s) = \begin{cases} 0 & \text{if } a = \epsilon \\ \text{Alt}(s) & \text{if } 0 = a \neq \epsilon \\ \text{Alt}(1 - s) & \text{if } 1 = a \neq \epsilon \end{cases}$$

Remind that 0 is the empty cycle and not zero as a function. Now the case $n > 2$:

$$\begin{aligned} \rho_1^\epsilon(A) &= \text{Alt} \sum_{i=1}^{n-1} \rho_1^\epsilon(A_i|A'_i) - \rho_1^\epsilon(A'_i|A_i) \\ \rho_1^\epsilon(A_i|A'_i) &= \left(\frac{s-t}{\epsilon-t}, \rho_1^1(A_i)(t), \rho_1^0(A'_i)(t) \right) + \delta_i \left(\frac{t-s}{\epsilon-t}, \rho_1^1(A_i)(t), \rho_1^0(A'_i)(s) \right) \\ &\quad + \delta_i \left(\frac{t-s}{\epsilon-t}, \rho_1^0(A_i)(t), \rho_1^1(A'_i)(s) \right) \end{aligned}$$

where $\delta_i = -\frac{1}{2}$ if $i \neq 1$ and $n-1$ and $\delta_1 = 0$, $\delta_{n-1} = -1$ (remind that $n > 2$). Now for $A = (a_1, \dots, a_n)$ let:

$$\begin{aligned} \rho_1(A)(s) &= \rho_1^0(A)(s) + \rho_1^1(A)(s) \\ \rho_k(A)(s_1, \dots, s_k) &= \sum_{i=1}^{n-k+1} \rho_1(A_i)(s_1) \rho_{k-1}(A'_i)(s_2, \dots, s_k). \end{aligned}$$

Proposition 3.9. *The morphisms ρ_k defined above, form a permuting integration theory on the binary subalgebra $\mathcal{C}_F^2$, generated by $(a_0; a_1, \dots, a_n; a_{n+1})$ with $a_i \in \{0, 1\}$ for $i = 1, \dots, n$.*

Remark 3.10. One can easily modify this construction to work for binary elements $(a_0; \dots; a_{n+1})$ with $a_i \in \{\alpha_0, \alpha_1\}$ for two fixed distinct elements $\alpha_0, \alpha_1 \in F$. The starting point would be:

$$\rho_1(a)(s) = \begin{cases} \frac{s-a}{\alpha_0-a} & \text{if } a \neq \alpha_0 \\ \frac{s-a}{\alpha_1-a} & \text{if } a = \alpha_0 \end{cases}$$

Now similarly we define ρ_1^0 and ρ_1^1 . Where instead of ϵ , we have α_ϵ .

Proof. According to lemma 3.7 all we need to do is to check ρ_1 is admissible and satisfies

$$d\rho_1 = -\delta\rho_2 - \sum_{\text{2-cuts of } A} \rho_1(A') \delta\rho_1(A'')$$

The admissibility follows from the following observation:

The intersection $\mathcal{F} \cap \rho_1^\epsilon(A)(s)$ of ρ_1^ϵ with any face of $\square^{2n-1}$ is empty if we set $s = \epsilon$. Note that this is stronger than just saying $\rho_1^\epsilon(\epsilon)$ is empty, which in fact is obvious since the first coordinate becomes 1. this stronger version follows since each component has both ρ_1^0 and ρ_1^1 , if we intersect with $z_1 = 0$ to kill the obvious coordinate giving 1 we still have persisting coordinates that gives 1, now regardless of ϵ . If we intersect with $z_1 = \infty$ we get the two terms $\rho_1^1(A_i)(\epsilon) \rho_1^0(A'_i)(s)$ and $\rho_1^0(A_i)(\epsilon) \rho_1^1(A'_i)(s)$. Now if $\epsilon = 0$ the second component is empty by induction and now for the first component we can use inductive hypothesis to show any intersection with a face becomes empty if we set $s = 0$, the case $\epsilon = 1$ is similar. This will imply the admissibility.

Now again as before the definition can be restated in terms of rooted binary trees with leaves at $a_1, \dots, a_n$. Such a tree gives a partition of this set into A_i and A'_i (left and right child labels). And to define ρ_1 or ρ_1^ϵ we need this partition. In fact for a binary tree T with left and right subtrees L and R we define:

$$\begin{aligned} \rho_1^\epsilon(T) = & \left(\frac{s-t}{\epsilon-t}, \rho_1^1(L)(t), \rho_1^0(R)(t) \right) + \delta_i \left(\frac{t-s}{\epsilon-t}, \rho_1^1(L)(t), \rho_1^0(R)(s) \right) \\ & + \delta_i \left(\frac{t-s}{\epsilon-t}, \rho_1^0(L)(t), \rho_1^1(R)(s) \right) + \left(\frac{s-t}{\epsilon-t}, \rho_1^0(L)(t), \rho_1^1(R)(t) \right) \\ & + \delta_{n-i} \left(\frac{t-s}{\epsilon-t}, \rho_1^0(L)(s), \rho_1^1(R)(t) \right) + \delta_{n-i} \left(\frac{t-s}{\epsilon-t}, \rho_1^1(L)(s), \rho_1^0(R)(t) \right) \end{aligned}$$

with δ_i as before with i being the number of leaves of R . With this definition we can write:

$$\rho_1^\epsilon(A) = \sum \rho_1^\epsilon(T)$$

where we sum over all binary trees with leaves labeled with $A = (a_1, \dots, a_n)$. Now take an edge e of one of these binary trees T , it corresponds to a coordinate z_e of $\rho_1^\epsilon(A)$. If it is not a leaf, in 6 different ways i.e. one of the 6 terms above. If one or both subtrees of this edge are leaves some of these 6 cases are empty. To manage these different choices we put an extra decoration on our binary trees. Namely we decorate each edge (excluding the leaves) with a number 1 through 6, corresponding to the above 6 terms in the order they are written. Now if (T, dec_T) is such a decorated tree with root decorated by $1 \leq j \leq 6$, we let $\rho_1^\epsilon(T, dec_T)$ be the j 'th term of the above expression, where L and R get their decorations from T . We can rewrite the above formula this time with decorated binary trees:

$$\rho_1^\epsilon(A) = \sum \rho_1^\epsilon(T, dec_T)$$

The 0-special edges of a decorated binary tree are the internal edge connected to the left of the root with the root decoration of 2 or 3 and the internal edge connected to the right of the root with the root decoration of 5 or 6 and the root itself. Similarly the ∞ -special edges are the internal ones connected to a leaf with a decoration of 2, 3, 5 or 6.

We want to show that there is an involution $(T, dec_T, e, \alpha) \leftrightarrow (T', dec_{T'}, e', \alpha')$ between quadruples a binary tree (with leaves at $a_1, \dots, a_n$), its decoration, an edge e and $\alpha = 0$ or ∞ such that with the right sign (dictated by the differential on cycles) $\rho_1^\epsilon(T, dec_T)|_{z_e=\alpha}$ cancels $\rho_1^\epsilon(T', dec_{T'})|_{z_{e'}=\alpha'}$, with the exclusion of (T, dec_T, e, α) where e is an α -special edge. This is done by considering different cases.

To calculate $d\rho^\epsilon(A)$ we only need to consider $\rho^\epsilon(T, dec_T)|_{z_e=\alpha}$ with e being an α -special edge for $\alpha = 0$ or ∞ . So we have the following cases. For ease of notation we right L^ϵ for $\rho_1^\epsilon(L, dec_L)(s)$ and similarly for R .

- (1) e is the root with decoration 1:

$$\rho_1^\epsilon(T, dec_T)|_{z_e=0} = L^1 R^0.$$

- (2) e is the root with decoration 4:

$$\rho_1^\epsilon(T, dec_T)|_{z_e=0} = L^0 R^1.$$

- (3) e is the root with decoration 2 or 6 (note that $\delta_i + \delta_{n-i} = 1$):

$$\rho_1^\epsilon(T, dec_T)|_{z_e=0} + \rho^\epsilon(T, dec'_T)|_{z_e=0} = -L^1 R^0.$$

where dec_T and dec'_T are identical except at the root, one with decoration 2 and one with decoration 6.

- (4) e is the root with decoration 3 or 5:

$$\rho_1^\epsilon(T, dec_T)|_{z_e=0} + \rho_1^\epsilon(T, dec'_T)|_{z_e=0} = -L^0 R^1.$$

- (5) e is to the left (or right) of the root with the root decoration 2 (or 6):

$$-\rho_1^0(T, dec_T)|_{z_e=0} + \rho_1^0(T, dec'_T)|_{z_e=0} = L^0 R^0.$$

$$-\rho_1^1(T, dec_T)|_{z_e=0} + \rho_1^1(T, dec'_T)|_{z_e=0} = -\delta_i L^1 R^0 - \delta_{n-i} L^0 R^1$$

- (6) e is to the left (or right) of the root with the root decoration 3 (or 5):

$$-\rho_1^1(T, dec_T)|_{z_e=0} + \rho_1^1(T, dec'_T)|_{z_e=0} = L^1 R^1$$

$$-\rho_1^0(T, dec_T)|_{z_e=0} + \rho_1^0(T, dec'_T)|_{z_e=0} = -\delta_i L^0 R^1 - \delta_{n-i} L^1 R^0$$

Adding all these terms will give us $L^0 R^0 + L^1 R^0 + L^1 R^1 + L^0 R^1$ or $(L^0 + L^1)(R^0 + R^1)$ when we sum over all decorated trees this gives $\rho_2(A)(s, s)$, which is a term in $-\delta\rho_2(A)$. The other terms in the formula:


$$d\rho_1(A) = -\delta\rho_2(A) + \sum \rho_1(A')\delta\rho_1(A'')$$

come from ∞ -special edges.

- (1) The edge e has a right child which is a leaf at a_{i+1} and a left child which is a tree T' with labels $A_i = (a_1, \dots, a_i)$, as shown below, with a decoration of 2 or 3 then:

$$\rho_1^\epsilon(T, dec_T)|_{z_e=\infty} + \rho_1^\epsilon(T, dec'_T)|_{z_e=\infty} = -\rho_1(T', dec_{T'})(a_{i+1})\rho_1^\epsilon(T'', dec_{T''})(s)$$

where T'' is the tree obtained from T by removing the leaf a_{i+1} and its sibling subtree T' and labeling the newly made leaf by a_{i+1} .




If we add the trivial term

$$-\rho_1(T', dec_{T'})(a_0)\rho_1^\epsilon(T'', dec_{T''})(s) + \rho_1(T', dec_{T'})(a_0)\rho_1^\epsilon(T'', dec_{T''})(s)$$

and add over all possible cases of this form we get the term

$$-\rho_2(A)(a_0, s) - \sum_i \rho_1(a_{i+1}, \dots, a_n)\delta\rho_1(a_0; a_1, \dots, a_i; a_{i+1})$$

- (2) The edge e has a left child which is a leaf at a_i and a right child which is a tree T' with labels $A'_i = (a_{i+1}, \dots, a_n)$ and decoration 4 or 5.




A calculation similar to the above case gives:

$$-\rho_2(A)(s, a_{n+1}) - \sum_i \rho_1(a_1, \dots, a_i) \delta \rho_1(a_i; a_{i+1}, \dots, a_n; a_{n+1})$$

- (3) One sees that the following two types of ∞ -special edges with $A_{i,j} = (a_{i+1}, \dots, a_j)$ will give the rest of the differential:

$$-\sum \rho_1(A') \delta \rho_1(A'')$$

where we sum over 2-cuts A', A'' of $(a_0; \dots; a_{n+1})$ such that the initial and final point of A'' are a_i and a_{j+1} with $0 < i < j < n$.



The proposition is proved. $\square$

In definition 2.12 we defined an ind-pro object $\Pi^C(S, a, b)$ for a given set $S \subseteq S'$ and a co-action:

$$\nu : \Pi^C(S, a, b) \longrightarrow \Pi^C(S, a, b) \otimes H^0 B(\tilde{\mathcal{C}}_{S, S'})$$

Now for $S = \{0, 1\}$ and $S' = \{0, 1, a, b\}$ we have a map from $\tilde{\mathcal{C}}_{S, S'}$ to $\mathcal{N}$ so we get a map

$$\begin{aligned} \Pi^C(S, a, b) &\longrightarrow \Pi^C(S, a, b) \otimes H^0 B(\mathcal{N}) \\ X_A &\mapsto \sum X_{A_1} \otimes \mathbb{I}_\rho(A_2) \cdots \mathbb{I}_\rho(A_k) \end{aligned}$$

where we sum over all elementary cuts of A . Which makes $\Pi^C(S, a, b) = \mathbb{Q}\langle\langle X_0, X_1 \rangle\rangle$ into a comodule over $H^0 B(\mathcal{N})$.

Definition 3.11. The motivic torsor of paths on $\mathbb{P}^1 - \{0, 1, \infty\}$ from a to b is the above co-module over $H^0 B(\mathcal{N})$. It is denoted by $\Pi^{Mot}(\mathbb{P}^1 - \{0, 1, \infty\}; a, b)$.

In the next section we show that its Hodge realization is the canonical Hodge structure on the unipotent completion of the torsor of paths. (see appendix B for a review of it)

4. HODGE REALIZATION OF THE MOTIVIC ITERATED INTEGRALS

As in definition 3.3, let $\{\rho_k\}$ be an integration theory, and $\mathbb{I}(A)$ its relative motivic iterated integral. In this section we show that the Hodge realization of $\mathbb{I}(A)$ is the canonical framed MHTS $\mathbb{I}^{\mathcal{H}}(A)$ (refer to definition B.7) for iterated integrals. Given $A = (a_0; a_1, \dots, a_n; a_{n+1})$, let $\mathcal{N}'$ be the sub algebra of $\mathcal{N}$ generated by $\rho(A_i)$'s for all cuts $A_1, \dots, A_k$ of A . We want to define a sub-DGA $\mathcal{D}'$ of topological cycles that satisfies the condition of the definition B.4 in appendix B. To do this we need to define some intermediate topological cycles. Their definition depends on a choice of a path γ with interior in $\mathbb{C} - \{a_1, \dots, a_n\}$ from the tangential base point a_0 to the tangential base point a_{n+1} .

For $1 \leq i \leq n$ define a topological cycles $\eta_i(A)$ inside $\square_{\mathbb{C}}^{2n-i}$ with (real) dimension $2n - i$ by:

$$\eta_i(A) = \rho_i(A)(\gamma(s_1), \dots, \gamma(s_i))$$

where s_k 's vary in the simplex $0 \leq s_1 \leq \dots \leq s_i \leq 1$.

Let Γ be a small disk around zero in $\square_{\mathbb{C}}$ with its canonical orientation, i.e. $\int_{\delta\Gamma} \frac{dz}{z} = 2\pi i$. We define:

$$\tau_i(A) = (-1)^i \delta \eta_i(A) \cdot \Gamma \cdot (\delta\Gamma)^{i-1} + \eta_i(A) \cdot (\delta\Gamma)^i$$

where $\cdot$ denotes the usual alternating product (i.e. we take the usual external product to obtain a cycle inside $\square^{2n}$ and then take the alteration with respect to the action of the group $\Sigma_{2n} \rtimes (\mathbb{Z}/2\mathbb{Z})^{2n}$). And δ denotes the topological boundary defined for a cycle $f(s_1, \dots, s_n)$ by:

$$\begin{aligned} (\delta f)(s_1, \dots, s_{n-1}) &= f(0, s_1, s_2, \dots, s_{n-1}) - f(s_1, s_1, s_2, \dots, s_{n-1}) + \dots \\ &\dots + (-1)^{n-1} f(s_1, \dots, s_{n-1}, s_{n-1}) + (-1)^n f(s_1, \dots, s_{n-1}, 1). \end{aligned}$$

Finally denote

$$\xi_{\gamma}(A) = \sum_{k=1}^n \tau_k(A).$$

Note that $\xi_{\gamma}(A) \in \tilde{\mathcal{D}}^0(n)$, i.e. it is a topological cycle inside $\square_{\mathbb{C}}^{2n}$ of (real) dimension $2n$. We will denote the corresponding element in the following homological version of $\tilde{\mathcal{D}}$ also by the same notation.

$$\mathcal{D}^0(n) = \text{Alt} \varinjlim H_{2n}(S \cup \mathcal{J}^{2n}, \mathcal{J}^{2n})$$

(where $\mathcal{J}^{2n}$ is the union of all the codimension 1 hyper planes of $(\mathbb{P}^1)^{2n}$ obtained by letting one coordinate equal to 1. The limit is taken over certain admissible subsets of $\mathbb{C}^{2n}$ of (real) dimension $2n$.)

We define $\mathcal{D}'$ to be the sub DGA of topological (homological) cycles generated by $\sigma(\rho(A_i))$ and $\xi_{\gamma}(A_i)$ where $A_1, \dots, A_n$ is any cut of A and $\sigma : \mathcal{N} \longrightarrow \mathcal{D}$ is the underlying topological cycle (i.e. the fundamental class). In fact one can take the algebra generated by these elements. It will automatically be a DGA because of the following proposition:

Proposition 4.1. *The differential of $\xi_{\gamma}(A)$ is given by the formula*

$$\sigma(\rho(A)) + \sum \xi_{\gamma}(A') \sigma(\rho(A''))$$

where we sum over all 2-cuts (A', A'') of A .

Proof. Since η_k were defined in terms of ρ_i 's, they satisfy:

$$\begin{aligned} d\eta_k(A) &= -\delta\eta_{k+1}(A) + (-1)^k \sum \eta_k(A')\sigma(\rho(A'')) \\ d(\delta\eta_k(A)) &= (-1)^k \sum \delta\eta_k(A')\sigma(\rho(A'')) \end{aligned}$$

Hence:

$$\begin{aligned} d\tau_k(A) &= (-1)^k d(\delta\eta_k(A) \cdot \Gamma \cdot \delta\Gamma^{k-1}) + d\eta_k(A) \cdot \delta\Gamma^k \\ &= (-1)^k d(\delta\eta_k(A)) \cdot \Gamma \cdot \delta\Gamma^{k-1} - \delta\eta_k(A) d(\Gamma) \delta\Gamma^{k-1} + d\eta_k(A) \cdot \delta\Gamma^k \\ &= \sum_{\text{2-cuts of } A} \left\{ \delta\eta_k(A')\sigma(\rho(A'')) \cdot \Gamma \cdot \delta\Gamma^{k-1} + (-1)^k \eta_k(A')\sigma(\rho(A'')) \cdot \delta\Gamma^k \right\} \\ &\quad + \delta\eta_k(A) \cdot \delta\Gamma^{k-1} - \delta\eta_{k+1}(A) \cdot \delta\Gamma^k \\ &= \sum_{\text{2-cuts of } A} \left\{ (-1)^k \delta\eta_k(A') \cdot \Gamma \cdot \delta\Gamma^{k-1} + \eta_k(A') \cdot \delta\Gamma^k \right\} \cdot \sigma(\rho(A'')) \\ &\quad + \delta\eta_k(A) \cdot \delta\Gamma^{k-1} - \delta\eta_{k+1}(A) \cdot \delta\Gamma^k \\ &= \delta\eta_k(A) \cdot \delta\Gamma^{k-1} - \delta\eta_{k+1}(A) \cdot \delta\Gamma^k + \sum_{\text{2-cuts of } A} \tau_k(A')\sigma(\rho(A'')) \end{aligned}$$

Summing over k we get:

$$d\xi_\gamma(A) = \delta\eta_1(A) + \sum_{\text{2-cuts of } A} \xi_\gamma(A') \cdot \sigma(\rho(A''))$$

but since $\delta\eta_1(A) = \sigma(\rho(A))$ the proposition is proved. $\square$

Finally our last piece of data, is a construction of certain element $Z_\gamma(A) \in H^0 B(\mathcal{D}', \mathcal{N}')$.

Definition 4.2. Define $Z_\gamma(A) \in H^0 B(\mathcal{D}', \mathcal{N}')^0$ by

$$\sum 1 \otimes [\rho(A_1)| \dots | \rho(A_k)] + \sum \xi_\gamma(A_1)[\rho(A_2)| \dots | \rho(A_k)]$$

where the sums are over all cuts of A .

Lemma 4.3. $dZ_\gamma(A) = 0$, hence it defines an element of $H^0 B(\mathcal{D}', \mathcal{N}')$ denoted by the same notation.

Proof. The first sum in Z_γ is in fact $1 \otimes \mathbb{I}(A)$. Since $d\mathbb{I}(A) = 0$ as an element of $B(\mathcal{N}')^0$, the only contribution of the differential for this term is

$$- \sum \sigma(\rho(A_1)) \otimes [\rho(A_2)| \dots | \rho(A_k)].$$

The external differential of each summand of the second sum is

$$d(\xi_\gamma(A_1)) \otimes [\rho(A_2)| \dots | \rho(A_k)] - \sum_{i=2}^n \xi_\gamma(A) \otimes [\rho(A_1)| \dots | d\rho(A_i)| \dots | \rho(A_k)]$$

Since $d\xi_\gamma(A_1) = \sigma(\rho(A_1)) + \sum \xi_\gamma(A'_1)\sigma(\rho(A''_1))$ and $d\rho(A_i) = -\sum \rho(A'_i)\rho(A''_i)$, adding the above two terms gives us:

$$\begin{aligned} &\sum_{\text{2-cuts of } A_1} \xi_\gamma(A'_1)\sigma(\rho(A''_1)) \otimes [\rho(A_2)| \dots | \rho(A_k)] \\ &+ \sum_{i=2}^n \xi_\gamma(A) \otimes [\rho(A_1)| \dots | \sum \rho(A'_i)\rho(A''_i)| \dots | \rho(A_k)] \end{aligned}$$

If we only sum this over k -cuts of A , it will be canceled by the internal differential of

$$\sum_{(k+1)\text{-cuts}} \xi_\gamma(A_1) \otimes [\rho(A_2)] \cdots [\rho(A_{k+1})].$$

□

Lemma 4.4. *Assume $a_0 \neq a_1$ and $a_n \neq a_{n+1}$. Under the map (see lemma B.5)*

$$\Lambda : H^0 B(\mathcal{D}', \mathcal{N}') \longrightarrow H^0 B(\mathcal{N}') \otimes \mathbb{C}$$

$Z_\gamma(A)$ maps to

$$\mathbb{I}(A) + \sum \left(\int_\gamma \omega_{A_1} \right) [\rho(A_2)] \cdots [\rho(A_k)]$$

where the sum is taken over all cuts $A_1, \dots, A_k$ of A , and if $A_1 = (a_0; a_{i_1}, \dots, a_{i_m}; a_{n+1})$

$$\omega_{A_1} = (2\pi i)^{-m} \frac{dt}{t - a_{i_1}} \circ \cdots \circ \frac{dt}{t - a_{i_m}}.$$

Proof. It is enough to show that

$$\lambda(\xi_\gamma(A_1)) = x^m \int_\gamma \omega_{A_1}.$$

where $m = \deg(A_1)$. To show this, first observe that if $i < m$

$$\int_{\eta_i(A_1)} \frac{dz_1}{z_1} \wedge \cdots \wedge \frac{dz_{2m-i}}{z_{2m-i}} = 0$$

This is so, since at least two components have the same parameter, and the alternation will kill them. By a similar argument we can, replace $\eta_i(A)$ with $\delta\eta_i$ and this time we can even include $i = m$. (This is the only place that one needs to look inside the machinery of integration theory). So the only term in $\xi_\gamma(A_1)$ that gives a nonzero contribution is $\eta_m(A_1) \cdot (\delta\Gamma)^m$. Since $\int_{\delta\Gamma} \frac{dz}{z} = 2\pi i$ we get:

$$\lambda(\xi_\gamma(A_1)) = x^m (2\pi i)^{-m} \int_{\eta_m(A_1)} \frac{dz_1}{z_1} \wedge \cdots \wedge \frac{dz_m}{z_m}.$$

It is left to check that:

$$\int_{\eta_m(A_1)} \frac{dz_1}{z_1} \wedge \cdots \wedge \frac{dz_m}{z_m} = \int_\gamma \frac{dt}{t - a_{i_1}} \circ \cdots \circ \frac{dt}{t - a_{i_m}}$$

But $\eta_m(A_1)$ is the cycle

$$\text{Alt}(c_1(\gamma(s_1) - a_{i_1}), \dots, c_m(\gamma(s_m) - a_{i_m}))$$

where $0 \leq s_1 \leq \dots \leq s_m \leq 1$ and the constants c_i depend on our integration theory. However since the form $\frac{dz_1}{z_1} \wedge \cdots \wedge \frac{dz_m}{z_m}$ is invariant under the action of $(\mathbb{C}^*)^m$ and is alternating under the action of $(\mathbb{Z}/2)^m \rtimes \Sigma_m$, we get the iterated integral on the right, by definition.

Finally let us show that the formula for $\Lambda(Z_\gamma(A))$ which a priori is in $B(\mathcal{N}) \otimes \mathbb{C}$, is in fact inside $H^0 B(\mathcal{N}) \otimes \mathbb{C}$. This is not needed because Λ is map of differential graded vector spaces, however it is not hard to see it directly. In fact if one fixes $A_1 = (a_0; a_{i_1}, \dots, a_{i_m}; a_{n+1})$ then $A_2, \dots, A_k$ are a shuffle of cuts for $(a_0; \dots; a_{i_1}), (a_{i_1}; \dots; a_{i_2}), \dots, (a_{i_m}; \dots; a_{n+1})$ and so the formula can be written as:

$$\mathbb{I}(A) + \sum \left(\int_\gamma \omega_{A_1} \right) \mathbb{I}(A_2) \cdots \mathbb{I}(A_k)$$

where this time we only sum over elementary cuts of A . $\square$

Lemma 4.5. *If $a_0 \neq a_1$ and $a_n \neq a_{n+1}$, the pair $(\mathcal{N}', \mathcal{D}')$ satisfies the conditions in definition B.4 of appendix B.*

Proof. Condition (1) follows from lemma 2.5. Condition (2) is trivial by definitions. Condition (3) follows from the calculations of the previous lemma. Conditions (4) and (6) are left to the reader to verify. To check condition (5) we need to show that $\tau^*(\mathbb{I}(A_i)) = 0$ where A_i is a cut component of A . However:

$$\mathbb{Z}_\gamma(A_i) = 1 \otimes \mathbb{I}(A_i) + \sum \xi_\gamma(A_{i1})[\rho(A_{i2}) | \dots | \rho(A_{ik})]$$

(summing over all cuts of A_i) has differential zero and $d(1 \otimes \mathbb{I}(A_i)) = -\tau(\mathbb{I}(A_i))$ so

$$\tau(\mathbb{I}(A_i)) = d(\sum \xi_\gamma(A_{i1})[\rho(A_{i2}) | \dots | \rho(A_{ik})]).$$

$\square$

Theorem 4.6. *Let $A = (a_0; a_1, \dots, a_n; a_{n+1})$. The Hodge realization of $\mathbb{I}(A)$ is $\mathbb{I}^{\mathcal{H}}(A)$.*

Proof. According to the lemma 2.11 for a permuting integration theory we can assume, $a_0 \neq a_1$ and $a_n \neq a_{n+1}$. Recall the MHTS's $\Pi^n(\mathbb{A}^1 - \{a_1, \dots, a_n\}; a_0, a_{n+1})$ and $H(\mathcal{N}', n)[n]$ of appendix B. Their $(-2k)$ -de Rham graded pieces are homogeneous degree k elements of $\mathbb{Q}\langle\langle X_{a_i} \rangle\rangle$ and $H^0 B(\mathcal{N}')(n-k)$ respectively. We define the following map from the first space to the second:

$$X_{b_1} \dots X_{b_k} \mapsto \sum [\rho(A_2) | \dots | \rho(A_m)]$$

where we sum over all cuts $A_1, \dots, A_m$ of A such that $A_1 = (a_0; b_1, \dots, b_k; a_{n+1})$. If there are no such cuts the value is zero and if $m = 1$, the value is 1. Using the same ideas as in the lemma 4.4, one can see that this in fact lands in $H^0 B(\mathcal{N}')(n-k)$. It maps 1 to $\mathbb{I}(A)$. It also maps $X_{a_1} \dots X_{a_n}$ to 1, while mapping any other monomial of degree n to zero. Hence it does respect the frames of $\mathbb{I}^{\mathcal{H}}(A)$ and $\text{Real}_{MHTS}(\mathbb{I}(A))$. It remains to show that it is a morphism of MHTS's. i.e. it maps the Betti subspace of $\mathbb{C}\langle\langle X_{a_i} \rangle\rangle$ defined by the image of

$$\Phi : \Pi^N(X, a_0, a_{n+1}) \longrightarrow \mathbb{C}\langle\langle X_{a_i} \rangle\rangle$$

$$\gamma \mapsto 1 + \sum_{b_i \in \{a_1, \dots, a_n\}} (2\pi i)^{-k} \left(\int_\gamma \frac{dt}{t - b_1} \circ \dots \circ \frac{dt}{t - b_k} \right) X_{b_1} \dots X_{b_k}$$

to the Betti subspace of $H^0 B(\mathcal{N}') \otimes \mathbb{C}$ defined by the image of the map Λ of lemma B.5. Observe that under the map defined between the two structures above, $\Phi(\gamma)$ maps to

$$\mathbb{I}(A) + \sum \left(\int_\gamma \omega_{A_1} \right) [\rho(A_2) | \dots | \rho(A_m)].$$

where we sum over all cuts of A . This is $\Lambda(Z_\gamma(A))$ and therefore is inside the Betti subspace of $H(\mathcal{N}', n)$. $\square$

APPENDIX A. BAR CONSTRUCTION

In this section we recall the basic definitions of bar construction. We follow the exposition of Hain and Zucker [HZ]. It differs from [BK] in certain signs.

A differential graded algebra (DGA for short) over a field F is a graded vector space over F :

$$\mathcal{A} = \bigoplus_{n \in \mathbb{Z}} \mathcal{A}^n$$

with an associative multiplication and linear maps:

$$d : \mathcal{A}^n \longrightarrow \mathcal{A}^{n+1}$$

such that $d \circ d = 0$. For $a \in \mathcal{A}^n$ and $b \in \mathcal{A}^m$, $a \cdot b \in \mathcal{A}^{n+m}$ and:

$$d(a \cdot b) = (da) \cdot b + (-1)^n a \cdot (db).$$

An augmentation for $\mathcal{A}$ is an onto map $\epsilon : \mathcal{A} \longrightarrow F$ of DGA's, where F is the trivial DGA, concentrated in degree 0. If we assume that $\mathcal{A}$ has a unit, we also require that $\epsilon(1) = 1$.

A (right) module over $\mathcal{A}$ is a graded vector space $\mathcal{M} = \bigoplus_n \mathcal{M}^n$ with a differential $d : \mathcal{M}^n \longrightarrow \mathcal{M}^{n+1}$ and an action:

$$\mathcal{M} \otimes \mathcal{A} \longrightarrow \mathcal{M}$$

such that:

- (1) For $m \in \mathcal{M}$ and $a, b \in \mathcal{A}$, $(ma)b = m(ab)$.
- (2) If $\mathcal{A}$ has a unit the $m \cdot 1 = m$.
- (3) For $m \in \mathcal{M}^n$ and $a \in \mathcal{A}^m$, $am \in \mathcal{M}^{n+m}$ and

$$d(ma) = (dm)a + (-1)^n m(da).$$

One can similarly define a left $\mathcal{A}$ module. We now give the details of the bar construction. Let $\mathcal{A}$ be an augmented DGA and $\mathcal{A}^+$ be the kernel of the augmentation. Let $\mathcal{M}$ be a right $\mathcal{A}$ module. The bar construction assigns a differential graded vector space $B(\mathcal{M}, \mathcal{A})$ to this data as follows. As a graded vector space it is:

$$\mathcal{M} \otimes_F T(\mathcal{A}^+[1]).$$

Here $T(\mathcal{A}^+[1])$ is the tensor algebra over the shifted graded vector space $\mathcal{A}^+[1]$. Hence the degree of a homogeneous element $m \otimes a_1 \otimes \dots \otimes a_r$ is

$$\deg(m) + \deg(a_1) + \dots + \deg(a_r) - r.$$

Often we use the bar notation: $m \otimes [a_1 | \dots | a_r]$ for such an element. There are two differential on this space, the external differential denoted by d_{ext} and the internal differential denoted by d_{int} . They are defined by the following formulae:

$$\begin{aligned} d_{ext}(m \otimes [a_1 | \dots | a_r]) &= dm \otimes [a_1 | \dots | a_r] \\ &\quad + \sum_{i=1}^r Jm \otimes [Ja_1 | \dots | Ja_{i-1} | da_i | a_{i+1} | \dots | a_r], \\ d_{int}(m \otimes [a_1 | \dots | a_r]) &= (Jm \cdot a_1) \otimes [a_2 | \dots | a_r] \\ &\quad + \sum_{i=1}^{r-1} Jm \otimes [Ja_1 | \dots | Ja_{i-1} | Ja_i \cdot a_{i+1} | \dots | a_r]. \end{aligned}$$

the operation J is given by $J(a) = (-1)^{\deg(a)-1}a$ on the homogeneous elements. Note that $\deg(a) - 1$ is the degree of a in the shifted complex $\mathcal{A}[1]$. If $\mathcal{M} = F$ is the trivial DGA, instead of $B(F, \mathcal{A})$ we write $B(\mathcal{A})$ and we write $[a_1] \dots [a_k]$ for $1 \otimes [a_1] \dots [a_k]$.

There is a coproduct on $B(\mathcal{A})$ (where the empty tensor is 1 by convention):

$$\Delta([a_1] \dots [a_r]) = \sum_{s=0}^r [a_1] \dots [a_s] \otimes [a_{s+1}] \dots [a_r].$$

Up to this point all the constructions work for a DGA which is not necessarily commutative. The product is defined only when both $\mathcal{M}$ and $\mathcal{A}$ are commutative DGA's (i.e. $ab = (-1)^{nm}ba$ for $a \in \mathcal{A}^n$ and $b \in \mathcal{A}^m$) by the shuffle:

$$m \otimes [a_1] \dots [a_r] \cdot m' \otimes [a_{r+1}] \dots [a_{r+s}] := \sum sgn_{\mathbf{a}}(\sigma)(m \cdot m') \otimes [a_{\sigma(1)}] \dots [a_{\sigma(r+s)}]$$

where σ runs over the (r, s) -shuffles and the sign is obtained by giving a_i 's weights $\deg(a_i) - 1$. For example if all a_i 's are of odd weights then all the signs are plus. These definitions make $B(\mathcal{A})$ into a Hopf algebra where $\mathcal{A}$ is a commutative DGA.

APPENDIX B. MIXED HODGE-TATE STRUCTURES

We recall a definition for the mixed Hodge-Tate structures (MHTS for short) below:

Definition B.1. A MHTS H is a finite dimensional $2\mathbb{Z}$ -graded $\mathbb{Q}$ -vector space $H_{dR} = \oplus_n H_{2n}$ together with a $\mathbb{Q}$ -subspace H_B of $H_{dR} \otimes \mathbb{C}$ such that³ for all m :

$$\text{Im} \left(H_B \cap \left(\bigoplus_{n \leq m} H_{2n} \otimes \mathbb{C} \right) \xrightarrow{Proj} H_{2m} \otimes \mathbb{C} \right) = H_{2m}.$$

A morphism f is a graded morphism f_{dR} from H_{dR} to H'_{dR} such that $f_{dR} \otimes 1 : H_{dR} \otimes \mathbb{C} \longrightarrow H'_{dR} \otimes \mathbb{C}$ sends H_B to H'_B , this induced map is denoted by f_B .

Remark B.2. If $H = (H_{dR}, H_B)$ is a MHTS as above, the shifted structure $H[n]$ is given by

$$H[n]_{dR} = \bigoplus_k H[n]_{2k}$$

$$H[n]_{2k} = H_{2k+2n}$$

and the same H_B .

Our two basic examples of such structures are the truncated torsor of paths on $X = \mathbb{A}^1 \setminus \{a_1, \dots, a_n\}$

$$\Pi^N(X; a_0, a_{n+1})$$

and the space $H^0 B(\mathcal{N}')$ for certain sub-DGA's $\mathcal{N}'$ of $\mathcal{N}$. Let us recall the construction of MHTS on these two spaces. For the first one we follow [G] §5 and for the second one we follow [BK] §8.

³It is more conventional to replace the right hand side with $(2\pi i)^{-m} H_{2m}$, however our definition is equivalent and make our constructions a little easier.

Assume that $a_1, \dots, a_n$ are distinct. Let H_{-2m} be the space of homogeneous polynomials of degree m on non-commuting variables $X_1, \dots, X_n$ with coefficients in $\mathbb{Q}$. This is a space of dimension m^n for $m > 0$ and 1 for $m = 0$ over $\mathbb{Q}$. Let

$$H_{dR} = \bigoplus_{0 \leq m \leq N} H_{-2m}.$$

Let $\omega_i := (2\pi i)^{-1} d \log(t - a_i)$. Any path $\gamma : [0, 1] \rightarrow X$ from a_0 to a_{n+1} defines an element of $H_{dR} \otimes \mathbb{C}$ by the Feynman-Dyson formula:

$$\Phi(\gamma) = \sum \left(\int_{\gamma} \omega_{i_1} \circ \dots \circ \omega_{i_k} \right) X_{i_1} \dots X_{i_k}$$

where we sum over all $(i_1, \dots, i_k)$ for $0 \leq k \leq N$ and $i_j \in \{1, \dots, n\}$. One can show that this element depends only on homotopy class of γ as a path from a_0 to a_{n+1} in X , and if we extend Φ by linearity to $\mathbb{Q}[\pi(X; a_0, a_{n+1})]$ it vanishes on $I_{a_0}^{N+1} \mathbb{Q}[\pi(X; a_0, a_{n+1})]$. I_{a_0} is the augmentation ideal $\text{Ker}(\mathbb{Q}[\pi_1(X; a_0)] \rightarrow \mathbb{Q})$. Hence we get a map

$$\Phi : \Pi^N(X; a_0, a_{n+1}) \rightarrow H_{dR} \otimes \mathbb{C}.$$

Its image is the Betti subspace H_B .

If a_0 or a_{n+1} are in $\{a_1, \dots, a_n\}$ one needs to consider tangential base point and regularization of the iterated integrals as follows. Identify the tangent space at each point of $\mathbb{C}$ with $\mathbb{C}$ and as the tangential base point at a , take the unit tangent vector 1. Now by a path from a_0 to a_{n+1} we mean a differentiable map $\gamma : [0, 1] \rightarrow \mathbb{C}$ such that $\gamma(0) = a_0$, $\gamma(1) = a_{n+1}$, $\gamma'(0) = \gamma'(1) = 1$ and γ restricted to the open interval $(0, 1)$ maps to $\mathbb{C} - \{a_1, \dots, a_n\}$. The regularized iterated integral is defined as follows. Let $\gamma^* \omega_i = f_i(t) dt$, then the integral

$$\int_{\epsilon \leq t_1 \leq \dots \leq t_k \leq 1-\epsilon} f_{i_1}(t_1) \dots f_{i_k}(t_k) dt_1 \dots dt_k$$

can be written as $\sum_{i=1}^m c_m(\epsilon) \log^i \epsilon$ where c_m are holomorphic around 0. The regularized value is $c_0(0)$.

Lemma B.3. *For a given N , the $2\mathbb{Z}$ -graded vector space*

$$H_{dR} = \bigoplus_{k \leq N} H_{-2k}$$

where H_{-2k} is the space of homogeneous polynomials of degree k on non-commuting variables $X_1, \dots, X_n$ with coefficients in $\mathbb{Q}$, together with the $\mathbb{Q}$ subspace H_B of $H_{dR} \otimes \mathbb{C}$ defined by:

$$\text{Im}(\Phi : \Pi^N(X; a_0, a_{n+1}) \rightarrow H_{dR} \otimes \mathbb{C})$$

is a MHTS. It is also denoted by $\Pi^N(X; a_0, a_{n+1})$.

Now we recall the MHTS defined in [BK] on $H^0 B(\mathcal{N}')$ for certain sub-DGA's $\mathcal{N}'$ of $\mathcal{N}$.

Let an admissible topological cycle of (real) codimension $2r$ in $\square_{\mathbb{C}}^n$ be a linear sum of admissible subsets of $\square_{\mathbb{C}}^n$ which are a union of smooth singular simplices of (real) codimension $2r$ and disjoint interiors. Admissible means that the intersection with any face of $\square^n$ is either empty or also a union of singular simplices of real codimension $2r$ and disjoint interiors. This group is denoted by $\text{TCycle}^r(n)$. Let

$$\tilde{\mathcal{D}}^n(r) = \text{Alt}(\text{TCycle}^r(2r - n) \otimes \mathbb{Q})$$

A similar constructions to $\mathcal{N}$, will make $\tilde{\mathcal{D}} = \oplus \tilde{\mathcal{D}}^n(r)$ into a DGA with a natural morphism:

$$\sigma : \mathcal{N} \longrightarrow \tilde{\mathcal{D}}$$

that assigns the underlying topological cycle to an algebraic cycle. We would like to consider these topological cycles up to homology. i.e. If for $Z \in \tilde{\mathcal{D}}^0(n)$ we have $dZ = 0$, we want it to be equivalent to $(\delta\Gamma)^{2n}$, where Γ is a small disk around zero in $\square_{\mathbb{C}}$ and δ is the boundary. This is done in more details in [BK]. But the idea is to consider

$$\mathcal{D}^n(r) := \text{Alt} \varinjlim H_{2r-2n}(S \cup \mathcal{J}^{2r-n}, \mathcal{J}^{2r-n})$$

where $\mathcal{J}^n$ is the union of all the codimension 1 hyper planes of $(\mathbb{P}^1)^n$ obtained by letting one coordinate equal to 1. The limit is taken over all admissible subset S in $\mathbb{C}^{2r-n}$ of (real) codimension $2r$. This has a natural structure of DGA (refer to [BK], §8).

Definition B.4. A pair of a sub-DGA $\mathcal{N}'$ of $\mathcal{N}$ and a sub-DGA $\mathcal{D}'$ of $\mathcal{D}$ is called admissible if:

- (1) Each space $H^0 B(\mathcal{N}')(r)$ is finite dimensional $\mathbb{Q}$ -vector space.
- (2) The map σ maps $\mathcal{N}'$ to $\mathcal{D}'$.
- (3) Let $\mathbb{C}[x]$ be a DGA concentrated in degree zero with Adams grading given by powers of x and zero differential. Define $\lambda : \mathcal{D}' \longrightarrow \mathbb{C}[x]$ to be zero on $(\mathcal{D}')^n(r)$ for $n \neq 0$ and for $c \in (\mathcal{D}')^0(r)$, i.e. an admissible topological cycle in $\square^{2r}$ of real dimension r :

$$\lambda(c) = \left((2\pi i)^{-2r} \int_c \frac{dz_1}{z_1} \wedge \dots \wedge \frac{dz_{2r}}{z_{2r}} \right) x^r$$

We will assume that λ is a well-defined morphism of DGA's. We call λ the period map.

- (4) The map $\lambda : H^0(\mathcal{D}') \longrightarrow H^0(\mathbb{C}[x]) = \mathbb{C}[x]$ defines an isomorphism $H^0(\mathcal{D}') = \mathbb{Q}[x]$. (This condition is missing from [BK], however is needed if one wants to prove that the following structure is a MHTS)
- (5) Let $\tau : B(\mathcal{N}') \rightarrow B(\mathcal{D}', \mathcal{N}')[1]$ (bracket denotes the shift in the degrees) be given by:

$$[a_1 | \dots | a_r] \mapsto \sigma(a_1) \otimes [a_2 | \dots | a_n].$$

It can easily be seen that this is a map of differential vector spaces so it induces:

$$\tau^* : H^0 B(\mathcal{N}') \longrightarrow H^1 B(\mathcal{D}', \mathcal{N}').$$

We will assume that this map is zero.

- (6) There is one more condition that is technical and we refer the reader to page 597 of [BK] for its statement.

Define

$$\Lambda : H^0 B(\mathcal{D}', \mathcal{N}') \longrightarrow H^0 B(\mathcal{N}') \otimes \mathbb{C}$$

via the period map followed by evaluation at $x = 1$: $\mathcal{D}' \xrightarrow{\lambda} \mathbb{C}[x] \xrightarrow{x=1} \mathbb{C}$.

Lemma B.5. For a given n , the $2\mathbb{Z}$ -graded vector space:

$$H_{dR} = \oplus_{r \leq n} H_{2r}, \quad H_{2r} = H^0 B(\mathcal{N}')(r)$$

together with the $\mathbb{Q}$ -subspace H_B of $H_{dR} \otimes \mathbb{C}$ given by:

$$\text{Im}(\Lambda : \oplus_{r \leq n} H^0 B(\mathcal{D}', \mathcal{N}')(r) \longrightarrow \oplus_{r \leq n} H^0 B(\mathcal{N}')(r) \otimes \mathbb{C})$$

is a MHTS. We denote this MHTS by $H(\mathcal{N}', n)$

The category of MHTS's is a neutral Tannakian category with a fiber functor given by sending H to H_{dR} . The general formalism of Tannakian categories imply that this category is equivalent to the category of graded comodules over a certain Hopf algebra. Beilinson observed that this Hopf algebra can be described using framed MHTS. An n -framed MHTS is a MHTS, H together with elements $v \in H_0$ and $\hat{v} \in \text{Hom}(H_{-2n}, \mathbb{Q})$. Two n -framed MHTS's, $(H, v, \hat{v})$ and $(H', v', \hat{v}')$ are equivalent if there is a morphism $f : H \longrightarrow H'$ such that $f_{dR,0}(v) = v'$ and $\hat{v}' \circ f_{dR,-2n} = \hat{v}$. This generates an equivalence relation, whose classes is denoted by $\chi_{MHTS}(n)$. The space $\chi_{MHTS} = \oplus_{n \geq 0} \chi_{MHTS}(n)$ has a natural structure of a Hopf algebra. The realization functor

$$\text{Real}_{MHTS} : H^0 B(\mathcal{N}) \longrightarrow \chi_{MHTS}$$

as a morphism of Hopf algebras has been defined in section 7 of [BK]. It is very complicated, however a more accessible definition at least for a sub-DGA, $\mathcal{N}'$ and a relative sub-DGA $\mathcal{D}'$ satisfying the above assumptions is also given.

Definition B.6. The Hodge realization map $\text{Real}_{MHTS} : H^0 B(\mathcal{N}') \longrightarrow \chi_{MHTS}$ is given by sending $a \in H^0 B(\mathcal{N}')(n)$ to the shifted MHTS $H(\mathcal{N}', n)[n]$ (ref. to the remark B.2) with the following frames:

$$\begin{aligned} a &\in H^0 B(\mathcal{N}')(n) = H(\mathcal{N}', n)[n]_0 \\ \epsilon &\in \text{Hom}(H^0 B(\mathcal{N}')(0), \mathbb{Q}) = \text{Hom}(H(\mathcal{N}', n)[n]_{-2n}, \mathbb{Q}) \end{aligned}$$

where ϵ is the augmentation.

Another important class of framed MHTS are associated to the iterated integrals.

Definition B.7. For a sequence $(a_0; a_1, \dots, a_n; a_{n+1})$ of elements of $\mathbb{C}$, the framed MHTS $\mathbb{I}^{\mathcal{H}}(a_0; a_1, \dots, a_n; a_{n+1})$ is defined by the MHTS $H = \Pi^n(\mathbb{C} - \{a_1, \dots, a_n\}; a_0, a_{n+1})$ with frames given by

$$\begin{aligned} 1 &\in H_0 = \mathbb{Q} \\ (X_{a_1} \cdots X_{a_n})' &\in \text{Hom}(H_{-2n}, \mathbb{Q}) = \text{Hom}(\mathbb{Q}\langle\langle X_{a_i} \rangle\rangle_{\deg=n}, \mathbb{Q}) \end{aligned}$$

where $(X_{a_1} \cdots X_{a_n})'$ is a morphism that sends $X_{a_1} \cdots X_{a_n}$ to 1 and any other monomial $X_{b_1} \cdots X_{b_n}$ with $b_i \in \{a_1, \dots, a_n\}$ to zero.

REFERENCES

- [BK] Bloch, Spencer; Křiž, Igor: Mixed Tate motives. Ann. of Math. (2) 140 (1994), no. 3, 557–605.
- [FJ] Furusho, Hidekazu; Jafari, Amir : Algebraic cycles and motivic generic iterated integrals. Math. research letters Vol. 14, No. 6, 2007, pp. 923 - 942.
- [GGL] Gangl, Herbert; Goncharov, Alexander B.; Levin, Andrei: Multiple logarithms, algebraic cycles and trees. Preprint. arXiv: math.NT 0504552.
- [G] Goncharov, Alexander B.: Multiple polylogarithms and mixed Tate motives. Preprint in math.AG/0103059.
- [HZ] Hain, Richard M.; Zucker, Steven: Unipotent variations of mixed Hodge structure. Invent. Math. 88 (1987), no. 1, 83124.